


ASTHMA ACTION PLAN

Take me when you visit your doctor

Photo (optional)

Name: _____
Plan date: _____ Review date: _____
Doctor details: _____

EMERGENCY CONTACT
Name: _____
Phone: _____
Relationship: _____


 **WELL CONTROLLED** is all of these...

- needing reliever medicine no more than 2 days/week
- no asthma at night
- no asthma when I wake up
- can do all my activities

Peak flow reading (if used) above _____

TAKE preventer
Name: _____
morning night puffs/inhalations
▪ Use my preventer, even when well controlled ▪ Use my spacer with my puffer

TAKE reliever
Name: _____
 puffs/inhalations as needed puffs/inhalations 15 minutes before exercise
▪ Always carry my reliever medicine

 **FLARE-UP** Asthma symptoms getting worse such as **any** of these...

- needing reliever medicine more than usual OR more than 2 days/week
- woke up overnight with asthma
- had asthma when I woke up
- can't do all my activities


Peak flow reading (if used) between _____ and _____
My triggers and symptoms

TAKE preventer
Name: _____
morning night puffs/inhalations for _____ days then back to **well controlled** dose

TAKE reliever
Name: _____ puffs/inhalations as needed

START other medicine
Name/dose/days/other treatments

MAKE appointment to see my doctor **same day or as soon as possible**

 **SEVERE** Asthma symptoms getting worse such as **any** of these...

- reliever medicine not lasting 3 hours
- woke up frequently overnight with asthma
- had asthma when I woke up
- difficulty breathing

Peak flow reading (if used) between _____ and _____
My triggers and symptoms


TAKE preventer
Name: _____
morning night puffs/inhalations for _____ days then back to **well controlled** dose

TAKE reliever
Name: _____ puffs/inhalations as needed

START other medicine
Name/dose/days/other treatments


MAKE appointment to see my doctor **TODAY**
▪ If unable to see my doctor, visit a hospital


OTHER INSTRUCTIONS
Other medicines, treatments, dose, duration, etc

 **EMERGENCY** is **any** of these...

- reliever medicine not working at all
- can't speak a full sentence
- extreme difficulty breathing
- feel asthma is out of control
- lips turning blue

Peak flow reading (if used) below _____

1  **CALL AMBULANCE NOW**
Dial Triple Zero (000)

2  **START ASTHMA FIRST AID**
Turn page for Asthma First Aid

ASTHMA FIRST AID

Blue/Grey Reliever

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- has a **known allergy to food, insects or medication and has SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available), even if there are no skin changes, then use a reliever**

1



SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
 - Repeat until 4 puffs have been taken



If using **Bricanyl**, give 2 separate inhalations (5 years or older)

If you don't have a spacer handy in an emergency, take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. **Repeat** until all puffs are given

3



WAIT 4 MINUTES

- If breathing does not return to normal, give 4 more separate puffs of reliever as above



Bricanyl: Give 1 more inhalation

IF BREATHING DOES NOT RETURN TO NORMAL

4



DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives



Bricanyl: Give 1 more inhalation every 4 minutes until emergency assistance arrives